

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 38/00</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 98/30227</b> <b>(43) International Publication Date:</b> 16 July 1998 (16.07.98)
<b>(21) International Application Number:</b> PCT/US98/00375 <b>(22) International Filing Date:</b> 12 January 1998 (12.01.98) <b>(30) Priority Data:</b> 119,989 10 January 1997 (10.01.97) IL <b>(71) Applicant (for all designated States except BB US):</b> YEDA RESEARCH AND DEVELOPMENT CO., LTD. [IL/IL]; P.O. Box 95, 76100 Rehovot (IL). <b>(71) Applicant (for BB only):</b> TEVA PHARMACEUTICALS USA CORP. [US/US]; 1510 Delp Drive, Kulpsville, PA 19443 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> ARNON, Ruth [IL/IL]; Weizmann Institute of Science, Meonot Shine 9, 76100 Rehovot (IL). SELA, Michael [IL/IL]; Weizmann Institute of Science, Meonot Wix 2, 76100 Rehovot (IL). TEIT-ELBAUM, Dvora [IL/IL]; Shkolnik Street 5, 76209 Rehovot (IL). GILBERT, Adrian [IL/IL]; Bialik Street 24/1, 44448 Kfar-Saba (IL). LINENBERG, Milka [IL/IL]; Sayfan Street 46, 40600 Tel-Mond (IL). RIVEN-KREITMAN, Rivka [IL/IL]; Dudaim Street 3/5, 44288 Kfar-Saba (IL).		<b>(74) Agents:</b> LEE, Steven, J. et al.; Kenyon & Kenyon, One Broadway, New York, NY 10004 (US).  <b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> TREATMENT OF MULTIPLE SCLEROSIS THROUGH INGESTION OR INHALATION OF COPOLYMER-1  <b>(57) Abstract</b>  The present invention relates to the treatment of multiple sclerosis by ingestion or inhalation of copolymer-1 and pharmaceutical compositions useful for such treatment.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## **TREATMENT OF MULTIPLE SCLEROSIS THROUGH INGESTION OR INHALATION OF COPOLYMER-1**

### **FIELD OF THE INVENTION**

This invention relates to the treatment of multiple sclerosis by ingestion or inhalation of copolymer-1 (as defined below). The present invention also relates to a pharmaceutical composition comprising copolymer-1 used for the treatment of multiple sclerosis, wherein the pharmaceutical composition is formulated for administration by ingestion or inhalation.

### **BACKGROUND OF THE INVENTION**

Copolymer-1, also known as glatiramer acetate and marketed under the tradename Copaxone®, comprises the acetate salts of polypeptides containing L-glutamic acid, L-alanine, L-tyrosine and L-lysine. The average molar fraction of the amino acids are 0.141, 0.427, 0.095 and 0.338, respectively, and the average molecular weight of copolymer-1 is between 4,700 and 11,000 daltons. It is a non-autoantigen which has been demonstrated to suppress experimental allergic encephalomyelitis (EAE) induced by various encephalitogens including mouse spinal cord homogenate (MSCH) which includes all myelin antigens, such as myelin basic protein (MBP) (Sela M et al., *Bull Inst Pasteur* (1990) 88 303-314), proteolipid protein (PLP) (Teitelbaum D et al., *J Neuroimmunol* (1996) 64 209-217) and myelin oligodendrocyte glycoprotein (MOG) (Ben-Nun A et al., *J Neurol* (1996) 243 (Suppl 1) S14-S22) in a variety of species. EAE is an accepted model for multiple sclerosis.

Copolymer-1 has been demonstrated to be active when injected subcutaneously, intra-peritoneally, intravenously or intramuscularly (D. Teitelbaum et al., *Eur.J.Immunol.* (1971) 1:242-248; D. Teitelbaum et al., *Eur.J.Immunol.* (1973) 3:273-279).

In phase III clinical trials, daily subcutaneous injections of copolymer-1

were found to slow progression of disability and reduce the relapse rate in exacerbating-remitting multiple sclerosis (K.P.Johnson, *Neurology* (1995) 1:65-70). Copolymer-1 therapy is presently limited to its daily subcutaneous administration.

5                   Currently, all specifically approved treatments of multiple sclerosis involve self-injection of the active substance. Frequently observed injection-site problems include irritation, hypersensitivity, inflammation, pain and even necrosis (in the case of at least one interferon  $\beta$  1-B treatment) and a low level of patient compliance. Therefore, an alternative method of administration is desirable.

10                   EP Patent 359,783 discloses the treatment of autoimmune diseases by oral administration of autoantigens. It discloses the oral administration of MBP for treatment of multiple sclerosis. Oral administration of an autoantigen has been termed "oral tolerance."

15                   PCT International Application Publication Nos. WO 91/12816, WO 91/08760, and WO 92/06704 all disclose the treatment of other autoimmune diseases using the "oral tolerance" method with a variety of autoantigens. However, none of these references disclose the treatment of multiple sclerosis by the oral  
20                   administration of non-autoantigen copolymer-1. The contents of all these patents and all literature references referred to above are hereby incorporated by reference in their entirety.

25                   It is, therefore, an object of the present invention to provide a method for treating multiple sclerosis by oral administration of copolymer-1 through ingestion or inhalation.

## BRIEF DESCRIPTION OF THE FIGURES

Figures 1 and 2 show the effect of copolymer-1 on the immune response to guinea pig myelin basic protein (GPBP) in rats (Figure 1) and mice (Figure 2) as assessed by spleen cell proliferation.

Figures 3 and 4 show the effect of copolymer-1 on cytokine release.

Figure 5 shows the suppression of EAE in mice by orally-administered copolymer-1. (SJL/JxBALB/c) F<sub>1</sub> mice were fed with PBS (■), 0.1 mg copolymer-1 (□), 0.25 mg copolymer-1 (▲), or 0.5mg copolymer-1 (Δ). Each dose was fed 7 times on days -7; -5; -3; 0; 2; 4 and 6. EAE was induced on day 0 by the injection of MSCH.

Figure 6 shows the proliferation and cytokine secretion by T-cell line derived from spleens of copolymer-1 fed rats. Cells were cultured with medium copolymer-1 (50 μ/ml) or concavalin A (Con A) (5 μg/ml). The proliferation and cytokine secretion responses to these antigens were measured.

Figure 7 shows the inhibition of EAE by T-cell line derived from spleens of copolymer-1 fed rats. Cells ( $20 \times 10^6$ /rat) were injected intraperitoneally 3 days after stimulation with copolymer-1, followed by EAE induction.

Figure 8 shows the inhibition of EAE by T-cell line derived from spleens of copolymer-1 fed mice. Cells ( $15 \times 10^6$ /mouse) were injected intravenously 3 days after stimulation with copolymer-1, followed by EAE induction.

FIG. 9 shows the Clinical Scores vs. Days of Copolymer-1 Post EAE Disease Induction in three Rhesus Monkeys.

FIG. 10 shows the Clinical Scores vs. Days of Copolymer-1 Post EAE

Disease Induction in six Rhesus Monkeys, in a trial comparing enteric-coated vs. uncoated pharmaceutical dosage form. Values at zero (0) have been separated on the y-axis to better show the results.

5

## DETAILED DESCRIPTION OF THE INVENTION

Accordingly, the present invention relates to the use of copolymer-1 in the preparation of a medicament for the treatment of multiple sclerosis. The medicament is administered through either ingestion or inhalation.

The present invention is also directed to a method of treating multiple sclerosis comprising administration of a therapeutically effective amount of copolymer-1, wherein the administration is through either ingestion or inhalation.

The present invention is further directed to a pharmaceutical composition administered through ingestion or inhalation comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of copolymer-1, wherein the pharmaceutical composition is used to treat multiple sclerosis.

As stated above, copolymer-1 comprises the acetate salts of polypeptides containing L-glutamic acid, L-alanine, L-tyrosine and L-lysine. The average molar fraction of the amino acids are 0.141, 0.427, 0.095 and 0.338, respectively, and the average molecular weight of copolymer-1 is between 4,700 and 11,000 daltons.

The present invention is based on the observation that, for example, the oral administration of copolymer-1 is effective in suppressing EAE, and, therefore, has a therapeutic value for the treatment of multiple sclerosis.

As contemplated, the copolymer-1 is brought into contact with those lymphoid tissues in the mucosal linings which are believed to be a primary source of immune system sensitization. These mucosal linings may be found (though not necessarily exclusively) in the sinuses, trachea, bronchial passages (where they are known as the BALT or bronchi-associated lymphoid tissues) and gastrointestinal linings (known as GALT or gut-associated lymphoid tissues). Thus, the administration of copolymer-1 is understood to include methods wherein copolymer-1 is introduced into the body by way of ingestion or inhalation. For example, copolymer-1 may be administered by way of the mouth through feeding, through a stomach tube, by inhalation into the bronchial passages or by nasal inhalation.

In one exemplary embodiment of the present invention, copolymer-1 is introduced orally in an amount of from 0.1 to 1000 mg per day, which may be administered as a single dose or in multiple dosages. As understood by one skilled in the art, the therapeutically effective dosage is generally a function of a patient's age, sex, and physical condition, as well as a function of other concurrent treatments being administered. The determination of the optimum, therapeutically effective dosage is well within the scope of one skilled in the art.

When copolymer-1 is introduced orally, it may be mixed with other food forms and consumed in solid, semi-solid, suspension, or emulsion form; and it may be mixed with pharmaceutically acceptable carriers, including water, suspending agents, emulsifying agents, flavor enhancers, and the like. In one embodiment, the oral composition is enterically-coated. Use of enteric coatings are well known in the art. For example, K. Lehman, *Acrylic Coatings in Controlled Release Tablet Manufacturer, Manufacturing Chemist and Aerosol News*, p. 39 (June 1973), and K. Lehman, *Programmed Drug Release From Oral Program Forms; Pharma. Int.*, vol. ISS 3 1971, p. 34-41, teach enteric coatings such as Eudragit S and Eudragit L. *Handbook of Pharmaceutical Excipients*, 2nd ed., also teaches Eudragit S and Eudragit L applications. One Eudragit which may be used in the present invention is L30D55.

Copolymer-1 may also be administered nasally in certain of the above-mentioned forms by inhalation or nose drops. Furthermore, oral inhalation may be employed to deliver copolymer-1 to the mucosal linings of the trachea and bronchial passages.

5

Copolymer-1 may be prepared by methods known in the art, for example, as disclosed in U.S. Patent No. 3,849,550, wherein the N-carboxyanhydrides of tyrosine, alanine,  $\gamma$ -benzyl glutamate and  $\epsilon$ -N-trifluoroacetyllysine are polymerized at ambient temperature in anhydrous dioxane with diethylamine as an initiator. The deblocking of the  $\gamma$ -carboxyl group of the glutamic acid is effected by hydrogen bromide in glacial acetic acid and is followed by the removal of the trifluoroacetyl groups from the lysine residues by 1M piperidine.

10

15

As described in PCT/WO95/31990, copolymer-1 having a desired average molecular weight of about  $7 \pm 2$  kilodaltons may preferably be prepared by a method comprising reacting protected copolymer-1 with hydrobromic acid to form trifluoroacetyl copolymer-1, treating the trifluoroacetyl copolymer-1 with an aqueous piperidine solution to form copolymer-1, and purifying the copolymer-1 so as to result in copolymer-1 having the desired average molecular weight.

20

The present invention will further be described in the examples below. However, the present invention should not be construed as being limited thereby. Unless otherwise indicated, all parts, percentages, and the like, are by weight.

25

#### EXAMPLE 1

30

Antigens - Copolymer-1, which was prepared according to the method described in PCT/WO95/31990, was obtained from Teva Pharmaceutical Industries Ltd., Israel. GPBP was prepared from guinea pig spinal cord by acid extraction and ammonium sulfate precipitation as described in Hirshfeld H et al., *Febs Lett* (1970)

7 317-320.

Animals - (PL/JxSJL/J) F1 female mice (8-10 weeks old) were obtained from Jackson Laboratories (Bar Harbor, ME). Female Lewis rats (8-12 weeks old) were obtained from Harlan-Olac (Bicester, G.B.).

Induction and Assessment of EAE - Mice were injected with 200 µg GPBP emulsified in an equal volume of complete Freund's adjuvant (CFA) containing 4 mg/ml mycobacterium tuberculosis (H37Ra) (Difco Lab, Detroit, Mich.). The emulsion at a total volume of 0.1 ml was injected into all four footpads. Immediately after and 24 hours later, pertussis toxin (250ng/mouse) (Sigma) was injected intravenously.

Rats were immunized with 25 µg of GPBP emulsified 1:1 in CFA containing 4 mg/ml H37Ra. The emulsion at a total volume of 0.1 ml. was injected into the two hindfoot pads.

Animals were examined daily from day 10 post induction for signs of disease. EAE was scored as follows: 0-no disease, 1-limp tail, 2-hind limb paralysis, 3-paralysis of all four limbs, 4-moribund condition, 5-death.

Induction of Oral Tolerance - Mice were fed with 250 µg GPBP or copolymer-1 dissolved in phosphate buffered saline (PBS) on days -7, -5, -3, 0, 2, 4 and 6 by gastric intubation with an 18-gauge stainless steel feeding needle (Thomas). EAE was induced on day 0.

Rats were fed with 1 mg GPBP or copolymer-1 dissolved in PBS by gastric intubation using a sterile feeding tube (Uno Plast, Denmark). Rats were fed 5 times (total dose of 5 mg) before disease induction at intervals of 2-3 days. EAE was induced two days after the last feeding. Control mice and rats were mock fed with PBS.

Proliferation Assay - The proliferation response of spleen cells was tested 10-11 days after EAE induction as described above. Cells from 3 animals in each group were pooled and cultured in triplicate ( $5 \times 10^5$  mouse cells and  $2 \times 10^5$  rat cells) in microtiter plates with various antigen concentrations (GPBP) in a final volume of 0.2ml. Microtiter plates contained RPMI 1640 (available from Sigma Biochemicals, St. Louis, Missouri) culture medium supplemented with 1% autologous serum. After 72 hr. of incubation, cells were pulsed with 1  $\mu$ Ci  $\{^3\text{H}\}$ -thymidine for 18 hr and then harvested onto filter papers and radioactivity was counted.

Cytokine Secretion Assay - Spleens were removed 10-11 days after EAE induction and cells of 3 mice from each group were pooled. Cells ( $5 \times 10^6$ /ml) were cultured in duplicates in 24 well plate in RPMI 1640 supplemented with 10% FCS (Fetal Calf Serum) in the presence or absence of antigen (GPBP 100 Fg/ml). Supernatants were harvested after 24-40 hr of culture. Quantitative ELISA for IL-2, IFN- $\gamma$ , IL-4, IL-6 and IL-10 were performed using paired mAbs specific for the corresponding cytokines (Pharmingen, La Jolla, CA) according to the manufacturer's instructions.

## Results:

The efficacy of orally administered copolymer-1 in preventing the clinical manifestations of EAE in Lewis rats was compared to that of GPBP, when assayed under conditions previously reported to induce oral suppression by GPBP (PJ Higgins & HL Weiner, *J Immunol* (1988) 140 440-445). The results summarized in Table 1 below demonstrate that copolymer-1 was more effective than GPBP and significantly decreased both the incidence (54% inhibition) and severity (57% inhibition) of EAE, as compared to PBS fed rats which served as control.

Table 1: Suppression of EAE in Rats by Oral Administration of Copolymer-1.

Fed Antigen	Incidence	Mean Maximal Score $\pm$ SD	Mean Onset (days)
PBS (Control)	27/28 (96%)	1.8 $\pm$ 0.5	11.9
GPBP	10/17 (59%) (p=0.0026)	0.9 $\pm$ 0.5	11.4
Copolymer-1	13/28 (46%) (p=0.00005)	0.78 $\pm$ 0.45	12.6

Each figure represents the cumulative results of 3-5 independent experiments. p values represent the statistical significance of difference from the control group (Fisher exact test). Mean maximal score was calculated for the entire group.

Effect of Antigen Feeding on the Immune Response to GPBP- The effect of oral administration of copolymer-1 and GPBP on the immune response to the disease inducing antigen-GPBP was tested in mice and rats. The results are summarized in Figure 1 which shows the reduction in cell proliferation by each of the orally administered compounds (copolymer-1 or GPBP) in a rat spleen cell suspension stimulated with GPBP (Figure 2 shows similar results from mice). As can be seen, oral administration of copolymer-1 resulted in almost complete inhibition of the proliferative responses to GPBP in these two species. In both species, copolymer-1 was more effective than GPBP in inhibiting the response to GPBP.

Cytokine levels were measured in the supernatants of spleen cell cultures derived from mice (Figures 3 and 4). Control mice fed with PBS secreted

IL-2; IFN- $\gamma$  and IL-6 (not shown) in response to GPBP. In mice fed with copolymer-1 or GPBP the amounts of the Th1 proinflammatory cytokines IL-2 and IFN- $\gamma$  produced in response to GPBP stimulation were lower than in control groups, with copolymer-1 being a more effective suppressant. IL-4 and IL-10 were not detected with any group treatment.

The results demonstrate that copolymer-1 is effective in suppressing EAE when given orally. The clinically protective effect of orally administered copolymer-1 is associated with down regulation of T cell immune responses to GPBP such as proliferation and proinflammatory cytokines (IL-2 and IFN- $\gamma$ ) release.

## EXAMPLE 2

Additional studies on the suppression of EAE by oral administration of copolymer-1 were performed in rats and mice. These studies established the optimal dose for treatment in each species. In order to understand the mechanism underlying oral suppression of EAE by copolymer-1, a copolymer-1 specific T-cell line was isolated from spleens of copolymer-1 fed animals. The *in vitro* reactivity of the lines and their *in vivo* effect on disease induction were studied.

### Materials and Methods:

Isolation of Copolymer-1 specific T cell lines - Lewis rats were fed 5 times with 1 mg copolymer-1 and (SJL/J x BALB/c) $F_1$  mice 7 times with 250  $\mu$ g copolymer-1, at intervals of 2-3 days. Four to twelve days after the last feeding animals were sacrificed and their spleens removed.

Spleen cells of 3 animals were pooled and incubated ( $50 \times 10^6$ /plate) with copolymer-1 (500  $\mu$ g) in medium containing 1% autologous serum for 4 days. Every 14-21 days cells ( $4-6 \times 10^6$ /plate) were restimulated by 3 days exposure to

copolymer-1 (500 µg) presented on syngeneic irradiated (3000 rad) rat thymocytes (100 x 10<sup>6</sup>/plate) or mouse splenocytes (50 x 10<sup>6</sup>/plate). Stimulation was followed by propagation in 10% supernatant of Con A activated normal mouse spleen cells as T cell growth factor (TCGF).

5

Proliferation Assay - T cell lines (1 x 10<sup>4</sup> cells) were cultured with irradiated (3000R) thymocytes (rat-1 x 10<sup>6</sup>) or splenocytes (mouse- 5 x 10<sup>5</sup>) and with the indicated antigens (10 µg copolymer-1; 1 µg Con A) in a final volume of 0.2 ml in microtiter plates. At the end of 48 hours incubation cultures were pulsed with <sup>3</sup>H-thymidine and harvested 6-12 hours later.

10

Cytokine Assay - T cells of rat line (0.5 x 10<sup>6</sup>/ml) were incubated with irradiated thymocytes (10 x 10<sup>6</sup>) with or without the indicated antigen (50µg copolymer-1, 5µg Con A). Cells were cultured for 24 hours in RPMI 1640 supplemented with 10% FCS for IL-2, TNFα, IL-4 and IL-10-measurements and in serum free medium - DCCM-1 (Biological Industries, Kibbutz Beit Haemek, Israel) for 72 hours for TGFβ measurement.

15

Cytokine levels in supernatants were measured in a quantitative ELISA using pairs of monoclonal antibodies specific for the corresponding cytokines.

20

Induction of EAE - (SJL/xBALB/c)F<sub>1</sub> mice were injected in all four footpads with 2mg mouse spinal cord homogenated (MSCH) emulsified in 1:1 ratio in CFA containing 1 mg/ml H37Ra (Difco Lab, Detroit, Mich). Pertussis toxin (250 ng/mouse, Sigma) was twice injected intravenously, once immediately after and again 48 hours later.

25

## RESULTS:

### 30 1. Dose response study in rats and mice

Rats were fed 5 times with 0.5, 1 or 2 mg copolymer-1 according to the established protocol (see Materials and Methods, above) and then challenged for EAE induction. The results are summarized in Table 2, and indicate that the most effective dose was 1 mg copolymer-1 – 0.5 mg or 2 mg copolymer-1 were less efficient in suppressing EAE.

(SJL/JxBALB/c) F<sub>1</sub> mice were fed 7 times with 0.1, 0.25 or 0.5 mg copolymer-1 on days -7; -5; -3; 0; 2; 4 and 6 by gastric intubation. EAE was induced on day 0 by the injection of MSCH. The results summarized in Figure 5 demonstrate that oral administration of copolymer-1 could suppress the disease in mice and the most effective dose was 0.1 mg copolymer-1. 0.25 mg of copolymer-1 was less effective and a dose of 0.5 mg was completely inactive. Thus, the results in both rats and mice demonstrate that oral copolymer-1 has an optimum dose response curve, and exceeding the effective oral dose resulted in inefficient suppression of EAE.

## 2. Studies with copolymer-1 specific Ts-lines established from copolymer-1 fed animals

Copolymer-1 specific T suppressor cell lines were isolated from spleens of rats and mice rendered unresponsive to EAE by feeding with copolymer-1. The proliferation and cytokine secretion response of such line isolated from rats is demonstrated in Figure 6. This line proliferated in response to copolymer-1 and secreted IL-2, some IL-10 and TGF $\beta$  but not TNF $\alpha$  or IL-4. This cytokine profile is compatible with Th3 type cells which were shown to be induced by oral MBP (Chen et al. Science 265, 1237, 1994).

The ability of the copolymer-1 specific lines to prevent EAE *in vivo* was studied. Cell lines were injected 3 days after *in vivo* stimulation with copolymer-1 (20 x 10<sup>6</sup> cells/rats injected i.p. and 15x10<sup>6</sup> cells/mouse injected i.v.). The animals were challenged for EAE induction immediately following cell transfer. The results illustrated in Figures 7 and 8 demonstrate that the disease was considerably inhibited

in the recipient animals. Thus, both the rat and murine copolymer-1 T-cell lines adoptively transferred the unresponsiveness to EAE induced by oral administration of copolymer-1. These cells actively downregulate the pathological immune response *in vivo*.

5

Table 2: Dose Response Study of Oral Copolymer-1 in Rats

	<u>Fed Antigen</u>	<u>Incidence</u>	<u>Mean Score ±SD</u>	<u>Mean Onset</u>	<u>Suppression* (%)</u>
10	PBS (control)	10/11	1.32±0.64	13.1	
	Copolymer-1 (0.5 mg)	9/11	0.95±0.57	13.5	28
15	Copolymer-1 (1 mg)	7/11	0.64±0.50	15.1	51
20	Copolymer-1 (2 mg)	8/11	0.91±0.70	13.1	31

\* calculated by mean score.

25 Each incidence figure represents the cumulative results of 2 individual experiments. Mean maximum clinical score was calculated for the whole group.

### Example 3

30

The effect of the oral administration of copolymer-1 on the induction of EAE in Rhesus monkeys was studied.

### Materials

35

Copolymer-1 was provided by Teva Pharmaceutical Industries Ltd. in an enteric-coated hard gelatin capsule comprising two dosage levels: 1 mg of copolymer-1 and 20 mg of copolymer-1. Each dosage level was formulated using mannitol, and coated with Eudragit L30D55. Placebo or control capsules comprised capsules containing 5 mg of sugar.

40

Bovine MBP was purchased from Life Technologies, Grand Island, New York. This material represents a highly purified preparation that gives a single band at 18.5 Kd following SDS-PAGE and Silver staining.

5     Feeding Protocol

3 Rhesus monkeys were treated as follows: One monkey served as control and was fed with placebo capsules (containing 5mg glucose only). The second and third monkeys were fed with copolymer-1-containing capsules at a dose of 1mg/feeding and 20mg/feeding, respectively. Animals were fed every other day for a total of 10 feedings: 5 times prior to disease induction (immunization on day 0) and then 5 times after immunization.

15     Disease Induction

Disease was induced on day 0 by an intradermal injection of 8mg bovine - MBP and 3 mg H37Ra *M. tuberculosis* in FCA into the hind footpads, total injected volume between 0.1 and 0.15 ml per footpad. The animals were followed for clinical manifestations of EAE, a variety of serum and CSF immunological markers and spinal cord and cranial MRI's.

20     Clinical Scoring

Symptom scores were given as follows: 0, normal neurological exam; 1, weight loss, anorexia, yawning, slow responses to stimuli, irritability or lethargy; 2, mild neurological signs, indifference, drooling, clumsiness using limbs, tremors, altered cry and disordered gaze; 3, moderate neurological signs, blindness (pupils do not react to light), akinesia, leg weakness, or paralysis; 4, severe neurological signs, semicoma, coma, quadriplegia; 5, death.

30

Antigen-induced lymphocytes proliferation

Heparinized blood samples were diluted 1:1 with Hanks balanced salt solution (BSS) containing 5% heat inactivated fetal calf serum (FCS), and layered in a hypaque-ficol gradient. Centrifugation (2000 rpm for 20 minutes at room temperature) allowed recovery of diluted plasma and the separation of lymphocytes at the interface. The recovered lymphocytes were washed three times in Hanks BSS-5% FCS and resuspended in RPMI 1640 complete medium containing RPMI 1640 medium, 10% FCS, and 1% of the following reagents: non-essential amino acids, sodium pyruvate, L-glutamine, 2-mercaptoethanol, and penicillin/streptomycin. The recovered lymphocytes were counted and resuspended at a final concentration of  $2 \times 10^6/\text{ml}$ . Cultures containing 100 microliters of the cell suspension and 100 microliters of MBP (20 micrograms/well), 100 microliters of copolymer-1 (10 micrograms/well) or 100 microliters of Con A (1 microgram/well) were set up in round bottom 96 well microtiter plates. The cultures were maintained at  $37^\circ\text{C}$  in 5%  $\text{CO}_2$  for 6 days. On day 5 of culture, each well was pulsed with  $1\mu\text{Ci}$  of  $^3\text{H}$ -thymidine for 16 to 18 hours. On day 6, the cultures were harvested by an automatic cell harvester, and counted by liquid scintillation methods. Stimulation indices were determined as follows:

$$\text{Stimulation index} = \frac{\text{Experimental cpm} - \text{background cpm}}{\text{background cpm}}$$

Clinical Results

No animal exhibited clinical symptoms for 24 days. The placebo-treated animal 18374 developed disease on day 25, and due to severe manifestations of EAE (score =  $4^+$  on a scale of 5) had to be sacrificed on day 28. The high-dose copolymer-1 treated animal 18498 treated with 20mg of copolymer-1 did not show any significant clinical symptoms over the 60 day observation period. The low-dose copolymer-1 treated animal 18497 began to show minimal symptoms on day 25,

however, in contrast with the placebo-treated animal 18374, '497 showed a much slower increase in clinical symptoms and leveled off at 2 to 3+ from day 28 to day 33. At this time-point animal '497 was fed with 20mg capsules 5 times on alternate days for a total of 10 days. As can be seen in Fig. 9, within 3 days the animal's clinical signs dropped to zero and remained there until the two copolymer-1 fed animals '497 and '498 were sacrificed on day 60 for histology.

#### Flow Cytometry

The Epics C flow cytometer was used to analyze both peripheral blood lymphocytes (PBL) and cerebral spinal fluid (CSF) collected from each monkey. The red blood cells were lysed and the remaining white blood cells (WBCs) were washed and stained using standard methods with appropriate reagents. Tables 3 and 4 below show the data from staining PBL and CSF WBCs from the three animals in this Example.

The number of cells staining CD4+ was slightly greater in copolymer-1 fed animals 18497 and 18498 than in the control animal 18374. The number of CD4+ cells that are also CD45RA- appears to increase in both control animal 18374 and low-dose copolymer-1 fed animal 18497 about the time both animals exhibited clinical symptoms (days +27 to +34), but remained fairly constant in the high-dose copolymer-1 fed animal 18498. The number of CD8+CD45RA+ staining cells decreased steadily in control animal 18374 and low-dose copolymer-1 fed animal 18497, but increased slightly in high-dose copolymer-1 fed animal 18498 (indicating the production of new CD4+ cells).

The number of cells found in the CSF of the control animal 18374 increased steadily from day 20+ until day 28+ when the animal died. Analysis showed that most of the CD4+ cells were CD45RA-. The number of cells collected from the CSF of animal 18497 were too few to count or stain. The number of cells collected from animal 18498 were too few to count or stain except on day +27. The CD4+ cells collected then were predominantly CD45RA-.

Table 3: Analysis of PBL obtained from Rhesus Monkeys immunized with MBP.

	Monkey	(Day of study)	CD4+	CD8+	Ratio	CD4+ CD45RA+	CD8+ CD45RA+
5	18374 (control)	(-1)	24	47	0.51	15	45
		(+6)	24	42	0.57	15	40
		(+13)	23	40	0.58	12	38
		+20	22	45	0.49	10	45
		+27	24	27	0.89	10	23
		+28	24	24	1.0	10	21
10	18497 (1mg Copolymer- 1/ dose)	-1	35	36	0.97	21	33
		+6	39	31	1.26	24	29
		+13	37	34	1.09	16	29
		+20	41	32	1.28	18	25
		+27	35	27	1.3	12	18
		+34	ND	ND	ND	ND	ND
20	18498 (20 mg Copolymer- 1/ dose)	-1	37	40	0.93	23	32
		+6	37	42	0.88	23	32
		+13	41	46	0.89	24	37
		+20	27	57	0.47	12	41
		+27	37	52	0.71	22	41
		+34	ND	ND	ND	ND	ND
25							

ND= not determined

Table 4: Analysis of CSF obtained from Rhesus Monkeys immunized with MBP.

Animal	(Day of study)	Cells $\mu$ l	CD4+	CD8+	CD4+:CD8+ Ratio	CD4+ CD45RA+	CD8+ CD45RA+
18374	0	ND	too few cells				
	+14	28	"				
	+20	100	61	18	2.18	2	4
	+28	294	35	35	0.94	4	8
18497	0	too few cells					
	+14	"					
	+20	"					
	+27	"					
	+34	"					
18498	0	too few cells					
	+14	"					
	+20	"					
	+27	29.7	47	ND		8	ND
	+34	too few cells					
	+41	"					

ND= not determined

Analysis For Antigen-specific T cell suppressor factor inducer (Tcsfi)

The analysis for MBP-specific Tsfi in the plasma of the three monkeys in this Example are seen in Tables 5 and 6 below. None of the animals produced MBP-specific Tsfi until day +13 following EAE induction with MBP. Control animal 18374 did not produce any Tsfi until day +20, and the level produced thereafter was just above background. Copolymer-1 fed animals 18497 and 18498 consistently produced significant levels of Tsfi from day +13 until day +41 shortly before termination.

Table 6 shows that plasma samples which did not contain Tsfi did

not react with anti-TGF-beta antibody. Plasma which exhibited MBP-binding with 3C9 antibody (anti-Tsfi) also reacted with the anti-TGF-beta antibody. Recombinant human TGF-beta reacted with the anti-TGF-beta, but not 3C9 antibody.

5

Table 5: Assay for MBP-specific T suppressor Inducer factor<sup>a</sup>

10

Animal #	-13	-1	+6	+13	+20	+27	+34	+41
18374	none	none	ND	ND	.17	.21	died	
18497	none	none	none	.60	.61	.56	.66	.76
18498	none	none	none	.32	.32	.42	.40	.40

15

a. Data represents OD at 405nm, of plasma samples at a 1:20 dilution  
ND = Not Determined

20

Table 6: The association between MBP-specific Tsfi and TGF-beta in plasma of Rhesus Monkeys treated with copolymer-1.

25

Animal #	Day of study	anti-Tsfi (3C9 Ab) <sup>a</sup>	anti-TGF-beta <sup>b</sup>
18374	-1	none	none
	+27	0.21	none
18497	+6	none	none
	+13	0.60	0.71
	+27	0.56	0.65
	+41	0.76	0.85
18498	+6	none	none
	+13	0.32	0.35
	+27	0.42	0.47
	+41	0.40	0.45
	r human TGF-beta <sup>c</sup>	-	0.72

30

- 35
- Represents OD405 nm of material bound for MBP (2.5µg of protein) and reacts with the anti-human Tsfi (3C9) antibody.
  - Represents OD405nm of material bound to MBP and reacts with anti-human TGF-beta antibody.
  - Represents well coated with 100ng of recombinant human TGF-beta.

40

Example 4

Six monkeys were treated and subsequent analyses performed substantially according to the protocol described for Example 3 hereinabove.

5

Control placebo-fed animal 18746 was fed with capsules containing glucose. Animal 18586 was fed with 1 mg copolymer-1 in capsules with cracked enteric coating. Animal 18639 was fed with 20 mg copolymer-1 in capsules with cracked enteric coating. Animal 18724 was fed with 1 mg copolymer-1 in intact enteric coated capsules. Animal 18810 was fed with 10 mg copolymer-1 in intact enteric coated capsules. Animal 18962 was fed with 20 mg copolymer-1 in intact enteric coated capsules.

10

The schedule of feeding, disease induction and follow-up were substantially the same as those of Example 3 described hereinabove.

15

Results

As can be seen with reference to Fig. 10, control monkey 18746 developed acute disease beginning on day +21 and died on day +23 of disease manifestations. Animals 18586 and 18639 treated with the "cracked" enteric coating, which opened in the stomach (at both doses), were not protected from the disease and died within 2-3 days of disease manifestations. All the monkeys fed with copolymer-1 in enteric coated capsule were fully protected from the disease, and developed no signs of EAE until day 60, when they were sacrificed for histology.

20

25

What is claimed is:

1. A method of treating multiple sclerosis comprising ingestion or inhalation of a therapeutically effective amount of copolymer-1.
2. A method according to claim 1, wherein the therapeutically effective amount of copolymer-1 is from 0.1 to 1000 mg/day.
3. A method in accordance with claim 1, wherein said copolymer-1 is administered orally.
4. A method in accordance with claim 1, wherein said copolymer-1 is administered nasally.
5. A method in accordance with claims 3 or 4, wherein said copolymer-1 is inhaled.
6. A method in accordance with claim 3, wherein said copolymer-1 is orally administered in an enterically-coated form.
7. The use of copolymer-1 for the manufacture of a medicament for the treatment of multiple sclerosis, wherein said medicament is administered through ingestion or inhalation.
8. The use as claimed in claim 7, wherein the medicament comprises from 0.1 to 1000 mg/day of copolymer-1.
9. The use according to claims 7 or 8, wherein the medicament is formulated for oral or nasal administration.
10. The use according to claim 9, wherein said administration is by inhalation.

11. The use according to claim 9, wherein the medicament is enterically-coated.
12. A pharmaceutical composition for the treatment of multiple sclerosis comprising a therapeutically effective amount of copolymer-1 and a pharmaceutically acceptable carrier, wherein said pharmaceutical composition is formulated for administration by either ingestion or inhalation.
13. The pharmaceutical composition of claim 12, wherein said pharmaceutical composition is in solid form, liquid form, aerosol or inhalable powder.
14. The pharmaceutical composition of claim 13, wherein said solid form is enterically-coated.
15. A method of treating multiple sclerosis in a subject comprising the administration of a therapeutically effective amount to the subject of the pharmaceutical composition claimed in any one of claims 12 to 14.

1/8




FIG. 2




FIG. 3





FIG. 4



Feeding Antigen

- PBS
- ▨ GPBP
- Cop1

3/8

FIGURE 5





FIGURE 6

# Proliferation and Cytokine Secretion by T-Cell Line Derived From Spleens of Cop1-Fed Rats.



FIGURE 7

## Inhibition of EAE by T-Cell Line Derived From Spleens of Cop1-Fed Rats



6/8

FIGURE 8

Inhibition of EAE by T cell Line Derived  
from Cop1-fed Mice






FIG 9

Clinical Score vs. Days of Cop-1 Post Disease Induction



Days After Immunization With MBP



## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/00375

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K 38/00

US CL :514/2

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/2

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Chem. abstr., Vol. 125, No. 23, 02 December 1996 (Columbus OH, USA), page 1, column 2, the abstract No. 291993b, JOHNSON, K.P., et al., 'Management of Relapsing/Remitting Multiple Sclerosis with Copolymer 1 (Copaxone).' Mult. Scler. 1996, 1(6), 325-326 (Eng), see entire document.	7-10, 12, 13 and 15
A		1-6, 11 and 14
A,P	US 5,719,296 A (ACTON, III et al.) 17 February 1998.	1-15

☐

Further documents are listed in the continuation of Box C.

☐

See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*B* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means	
*P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

17 FEBRUARY 1998

Date of mailing of the international search report

13 APR 1998

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

RAYMOND J. HENLEY III

Telephone No. (703) 308-1235